

SUPPLEMENTAL MATERIAL

Supplementary table 1: Diagnostic criteria for severity of PEP

	Consensus (3)	Revised Atlanta (4)
Mild	Hospitalization of 2-3 days	- No organ failure
Moderate	Hospitalization of 4-10 days	- No local or systemic complications
		- Organ failure that resolves within 48h (transient organ failure) and/or
		- Local or systemic complications without persistent organ failure
Severe	Hospitalization of >10 days, or hemorrhagic pancreatitis, phlegmon, or pseudocyst, or intervention (percutaneous drainage or surgery)*	- Persistent organ failure (>48h) <ul style="list-style-type: none"> o Single organ failure o Multiple organ failure

*Hemorrhagic pancreatitis was defined as necrotizing pancreatitis and the term 'phlegmon' was abandoned (14).

Supplementary table 2: Diagnostic criteria for PEP

Consensus	Revised Atlanta
- New or increased abdominal pain consistent with pancreatitis	- Abdominal pain
- Serum amylase/lipase ≥ 3 x ULN at more than 24 hours after ERCP	- Serum amylase/lipase >3 x ULN
- Admission or prolongation of planned admission for at least 2 days	- Acute pancreatitis on CE-CT, MRI or US

CE-CT: contrast enhanced computed tomography scan. ERCP: endoscopic retrograde cholangiopancreatography. MRI: magnetic resonance imaging. ULN: upper limit of normal. US: ultrasound.